

REFERENCES

- Algase, D., Beck, C., Kolanowski, A., Whall, A., Berent, S., Richards, K., and Beattie, E. (1996). Need-driven dementia-compromised behavior: An alternative view of disruptive behavior. *American Journal of Alzheimer's disease*, 11(6): 10-19.
- Allen-Burge, R., Stevens, A., and Burgio, L. (1999). Effective behavioral interventions for decreasing dementia-related challenging behavior in nursing homes. *International Journal of Geriatric Psychiatry*, 14: 213-232.
- Alzheimer's Association. (2000). *Issue Kit: Special Care Units*. Alzheimer's Association State Policy Clearinghouse.
- Archibald, C. (1999). Commentary on Effective behavioral interventions for decreasing dementia-related challenging behavior in nursing homes by Allen-Burge et al. *International Journal of Geriatric Psychiatry*, 14: 228-230.
- Banaszak-Holl, J., Zinn, J., and Mor, V. (1996). The impact of market and organizational characteristics on nursing care facility service innovation. A resource dependency perspective. *Health Services Research*, 31(1): 97-117.
- Beck, C., Frank, L., Chumbler, N., O'Sullivan, P., Vogelpohl, T., Rasin, J., Walls, R., and Baldwin, B. (1998). Correlates of disruptive behavior in severely cognitively impaired nursing home residents. *The Gerontologist*, 38(2): 189-198.
- Boucher, L. (1999). Disruptive behaviors in individuals with Alzheimer's disease: A behavioral approach. *American Journal of Alzheimer's Disease*, 14(6): 351-356.
- Braceland Center for Mental Health and Aging, email from Julie Robison, Ph.D., August 11, 2004.
- Brazil, K., Hasler, A., McAiney, C., Sturdy-Smith, C., and Tettman, M. (2003). Perceptions of resident behavior problems and their clinical management in long term care facilities. *Journal of Mental Health and Aging*, 9(1): 35-42.
- Bridges-Parlet, S., Knopman, D., and Thompson, T. (1994). A descriptive study of physically aggressive behavior in dementia by direct observation. *Journal of the American Geriatrics Society*, 42: 192-197.

- Brodaty, H., Draper, B., Saab, D., Lee-Fay, L., Richards, V., Paton, H., and Lie, D. (2001). Psychosis, depression and behavioral disturbances in Sydney nursing home residents: Prevalence and predictors. *International Journal of Geriatric Psychiatry*, 16: 5104-512.
- Brunk, D. (1997). Random acts of violence. *Contemporary Long Term Care*, July, 20(7), 38-49, 42.
- Burgio, L., Stevens, A., Burgio, K., Roth, D., Paul, P., and Gerstle, J. (2002). Teaching and maintaining behavior management skills in the nursing home. *The Gerontologist*, 42(4): 487-496.
- Burgio, L. and Stevens, A. (1999). Behavioral interventions and motivational systems in the nursing home. *Annual Review of Gerontology and Geriatrics*, 18: 284-320.
- Burgio, L. and Scilley, K. (1994). Caregiver performance in the nursing home: The use of staff training and management procedures. *Seminars in Speech and Language*, November, 15(4): 313-322.
- Burgio, L. and Bourgeois, M. (1992). Treating severe behavioral disorders in geriatric residential settings. *Behavioral Residential Treatment*, 7(2): 145-168.
- Burgio, L. and Burgio, K. (1990). Institutional staff training and management: A review of the literature and a model for geriatric, long-term care facilities. *International Journal of Aging and Human Development*, 30(4): 287-302.
- Camp, C., Cohen-Mansfield, J., and Capezuti, E. (2002). Use of nonpharmacologic interventions among nursing home residents with dementia. *Psychiatric Services*, November, 53(11): 1397-1401.
- Carlson, D., Fleming, K., Smith, G., and Evans, J. (1995). Management of dementia-related behavioral disturbances: A nonpharmacologic approach. *Mayo Clin Proc*, 70: 1108-1115.
- Castle, N. and Fogel, B. (1998). Organizational structure and outcomes for nursing home residents with mental disorders. *Journal of Mental Health and Aging*, 4(1): 105-124.
- Cawood, J. (2002). Mental illness and physical aggression: Solutions for long-term care facilities. *Long-Term Care Interface*, January: 26-34.

- Chartok, P., Nevins, A., Rzetelny, H., and Gilberto, P. (1988). A mental health training program in nursing homes. *The Gerontologist*, 28(4): 503-507.
- Class, C., Schneider, L., and Farlow, M. (1997). Optimal management of behavioral disorders associated with dementia. *Drugs and Aging*, 10(2): 95-106.
- Cody, M., Beck, C., and Svarstad, B. (2002). Challenges to the use of nonpharmacologic interventions in nursing homes. *Psychiatric Services*, 53(11): 1402-1406.
- Cohen-Mansfield, J. (1995). Assessment of disruptive behavior/agitation in the elderly: Function, methods, and difficulties. *Journal of Geriatric Psychiatry and Neurology*, 8: 52-60.
- Cohen-Mansfield, J., Marx, M., and Werner, P. (1992). Agitation in elderly persons: An integrative report on findings in a nursing home. *International Psychogeriatrics*, 4(suppl.2): 221-239.
- Distasio, C. (1994). Violence in health care: Institutional strategies to cope with the phenomenon. *Health Care Supervisor*, 12(4): 1-34.
- Eastman, P. (2004). Closer to a cure for Alzheimer's. *AARP Bulletin*, 45(7): 12-14.
- Emerson Lombardo, N., Fogel, B., Robinson, G., and Weiss, H. (1996). *Achieving Mental Health of Nursing Home Residents: Overcoming Barriers to Mental Health Care*. Hebrew Rehabilitation Center for Aged, Boston, MA and Mental Health Policy Resource Center, Washington, D.C.
- Emerson Lombardo, N. (1994). *Barriers to Mental Health Services for Nursing Home Residents*. AARP Public Policy Institute.
- Gerdner, L. and Buckwalter, K. (1996). Review of state policies regarding special care units: Implications for family consumers and health care professionals. *The American Journal of Alzheimer's Disease*, 11(2): 16-27.
- German, P., Rovner, B., Burton, L., Brant, L., and Clark, R. (1992). The role of morbidity in the nursing home experience. *The Gerontologist*, 32(2): 152-158.

- Hagen, B. and Sayers, D. (1995). When caring leaves bruises: The effects of staff education on resident aggression. *Journal of Gerontological Nursing*, 21(11): 7-16.
- Holmes, D., Teresi, J., Weiner, A., Monaco, C., Ronch, J., and Vickers, R. (1990). Impacts associated with special care units in long term care facilities. *The Gerontologist*, 30: 178-183.
- Hughes, T. and Medina-Walpole, A. (2000). Implementation of an interdisciplinary behavior management program. *Journal of American Geriatric Society*, 48: 581-587.
- Jackson, M., Spector, W., and Rabins, P. (1997). Risk of behavior problems among nursing home residents in the United States. *Journal of Aging and Health*, 9(4): 451-472.
- Kanapaux, W. (2004). Many needs but few psychiatric services for seniors in long-term care. *Psychiatric Times*, 23(13). Retrieved from the World Wide Web: [http://: www.psychiatrictimes.com/p041107.html](http://www.psychiatrictimes.com/p041107.html) on December 8, 2004.
- Kennedy, G. Poor delivery of geriatric mental health services in long-term care setting: Implications for the shaping of public policy and nursing home regulations. *Long-Term Care Forum: Care of the Psychiatric Patient in the Nursing Home: Challenges and Opportunities*, 1(3). Retrieved from the World Wide Web: <http://www.aagpga.org/prof/default.asp> on June 30, 2004.
- Kitwood, T. and Bredin, K. (1992). Towards a theory of dementia care: Personhood and well-being. *Ageing and Society*, 12: 269-287.
- Kolanowski, A., Litaker, M., and Baumann, M. (2002). Theory-based intervention for dementia behaviors: A within-person analysis over time. *Applied Nursing Research*, May, 15(2): 87-96.
- Kolanowski, A, Hurwitz, S, Taylor, L., Evans, L, and Strumpf, N. (1994). Contextual factors associated with disturbing behaviors in institutionalized elders. *Nursing Research*, 43(2): 73-79.
- Landreville, P., Bordes, M., Dicaire, L., and Verreault, R. (1998). Behavioral approaches for reducing agitation in residents of long-term care facilities: Critical review and suggestions for future research. *International Psychogeriatrics*, 10(4): 397-419.

- Lennox, D. (2002). Is your long-term care facility behaviorally prepared? *Healthcare Review*, July, 15(6), 5(1). Retrieved from iConn, the Connecticut Digital Library, Infotrac on May 10, 2004.
- Low, L., Draper, B., and Brodaty, H. (2004). The relationship between self-destructive behavior and nursing home environment. *Aging and Mental Health*, January, 8(1): 29-33.
- Lyketsos, Constantine L. A strategic approach to the psychiatric care of memory-impaired patients. *Long-Term Care Forum: Psychotherapeutic Management of the Long-Term Care Patient*, 1(2). Retrieved from the World Wide Web: <http://www.aagapga.org/prof/default.asp> on June 30, 2004.
- McGrew, K. (1999). Residents with severe mental illnesses: How nursing homes respond. *Journal of Gerontological Social Work*, 31(3/4): 149-168.
- Mor, V., Banaszak-Holl, J., and Zinn, J. (1995-1996). The trend toward specialization in nursing care facilities. *Generations*, Winter: 24-29.
- Morgan, D., and Stewart, N. (1997). The importance of the social environment in dementia care. *Western Journal of Nursing Research*, December, 10(6): 740(22). Retrieved from iConn, the Connecticut Digital Library, Infotrac on May 10, 2004.
- Morrison, E., Ramsey, A., and Synder, B. (2000). Managing the care of complex, difficult patients in the medical-surgical setting. *MEDSURG Nursing*, 9(1): 21-26.
- Morrison, E. (1994). The evolution of a concept: Aggression and violence in psychiatric settings. *Archives of Psychiatric Nursing*, 8(4): 245-253.
- Morrison, E. (1992). A coercive interactional style as an antecedent to aggression in psychiatric patients. *Research in Nursing and Health*, 15: 421-431.
- O'Brien, J., Shomphe, L., and Caro, J. (2000). Behavioral and psychological symptoms in dementia in nursing home residents: The economic implications. *International Psychogeriatrics*, 12(Suppl 1): 51-57.
- Occupational Safety and Health Administration, U.S. Department of Labor. (2004). *Guidelines for Preventing Workplace Violence for Health Care and Social Service Workers*. OSHA 3148-O1R.

Omnibus Budget Reconciliation Act, 1987, 42 CFR 483.10.

Opie, J., Doyle, C., and O'Connor, D. (2002). Challenging behaviors in nursing home residents with dementia: A randomized controlled trial of multidisciplinary interventions. *International Journal of Geriatric Psychiatry, 17*: 6-13.

Radebaugh, T., Buckholtz, N., and Khachaturian, Z. (1996). Behavioral approaches to the treatment of Alzheimer's disease: Research strategies. *International Psychogeriatrics, 8*(Suppl 1): 7-12.

Reichman, William E. Quality of psychiatric services in the nursing home: Impact on patients, staff, and the nursing home milieu. *Long-Term Care Forum: Care of the Psychiatric Patient in the Nursing Home: Challenges and Opportunities, 1*(3). Retrieved from the World Wide Web: <http://www.aagapga.org/prof/default.asp> on June 30, 2004.

Richter, J. (1989). Providing nursing home care to the chronically mentally ill. *Journal of Gerontological Nursing, 15*(6): 18-23.

Rohrer, J., Buckwalter, K., and Russell, D. (1989). The effects of mental dysfunction on nursing home care. *Soc Sci Med, 28*(4): 399-403.

Ryden, M., Bossenmaier, M., and McLachlan, C. (1991). Aggressive behavior in cognitively impaired nursing home residents. *Research in Nursing and Health, 14*: 87-95.

Schnelle, J., Ouslander, J., and Cruise, P. (1997). Policy without technology: A barrier to improving nursing home care. *The Gerontologist, 37*(4): 527-532.

Simonowitz, J. (1995). Violence in health care: A strategic approach. *Nurse Practitioner Forum, 6*(2): 120-129.

Sloane, P., Lindeman, D., Phillips, C., Morita, D., and Koch, D. (1995). Evaluating Alzheimer's special care units: Reviewing the evidence and identifying potential sources of study bias. *The Gerontologist, 35*(1): 103-111.

Spector, W. and Jackson, M. (1994). Correlates of disruptive behaviors in nursing homes. *Journal of Aging and Health, May, 6*(2): 173-184.

Spore, D., Smyer, M., and Cohn, M. (1991). Assessing nursing assistants' knowledge

of behavioral approaches to mental health problems. *The Gerontologist*, 31(3): 309-317.

State of Connecticut, General Assembly, *Final Report of the Ad Hoc Task Force on Nursing Home Costs in Connecticut*, February 15, 2002.

State of Connecticut, Department of Mental Health and Addiction Services. (2001). *Report on Violence in Nursing Homes*.

State of Connecticut, Department of Public Health, conversation with Diane Smith, June 3, 2004.

State of Connecticut, Legislative Program Review and Investigations Committee, *Staffing in Nursing Homes*. Retrieved from the World Wide Web: <http://www.cga.gov/pri/archives/2000sdigest.htm> on December 17, 2004.

State of Connecticut, Legislative Program Review and Investigations Committee, *Medicaid Rate Setting for Nursing Homes*. Retrieved from the World Wide Web: <http://www.cga.ct.gov/archives/2001nhdigest.htm> on December 17, 2004.

State of Connecticut, Office of Policy and Management Population Projections. Series 95.1, 1995.

Stevens, A., Burgio, L., Bailey, E., Burgio, K., Paul, P., Capilouto, E., Nicovich, P., and Hale, G. (1998). Teaching and maintaining behavior management skills with nursing assistants in a nursing home. *The Gerontologist*, 38(3): 379-384.

Stevenson, S. (1991). Heading off violence with verbal de-escalation. *Journal of Psychosocial Nursing*, 29(11): 6-10.

Sutor, B., Rummans, T., and Smith, G. (2001). Assessment and management of behavioral disturbances in nursing home patients with dementia. *Mayo Clin Proc*, 76: 540-550.

U.S. Department on Health and Human Services, Center for Medicare and Medicaid Services, MDS Reports retrieved from the World Wide Web: <http://www.cms.hhs.gov/states/mdsreports> on August 8, 2004.

U.S. Department on Health and Human Services, Center for Medicare and Medicaid Services Media Affairs Press Release, Wednesday, December 22, 2004. *Nursing Home Care Improving in Many Areas New CMS Data Show, New Steps Initiated.*

Van Haitsma, K., Lawton, M., and Kleban, M. (2000). Does segregation help or hinder?: Examining the role of homogeneity in behavioral and emotional aspects of quality of life for persons with cognitive impairment in the nursing home. *Research and Practice in Alzheimer's Disease*, 4: 163-177.

Watts, G. (2001). The development of a service for older people with challenging behavior. *Signpost*, 6(2): 38-39.

Whall, A. and Kolanowski, A. (2004). Editorial: The need-driven dementia-compromised behavior model - a framework for understanding the behavioral symptoms of dementia. *Aging and Mental Health*, March, 8(2): 106-108.